

Greenhouse and High Tunnel Coverings



Cary L. Rivard, Ph.D.

**Dept. of Horticulture and
Natural Resources**

K-STATE
Research and Extension

Pre-harvest effects on postharvest quality of fresh produce



Dr. Eleni Pliakoni
Urban Food Production
and Postharvest
Physiology

KSU Olathe







The Next Frontier in Food



- Standard poly will reduce PAR 7-10%
- “Diffuse” poly coverings
- Ability to implement shade cloth
- Poly removal..?





3D




Why Does it Matter

Plant Physiology 101: Photosynthesis




- Plants are constantly “quenching” electrons
- Carotenoids and anthocyanins

Plant Physiology 101: Pigments



Production of Organic Tomatoes in High Tunnels is reducing Pre-Harvest and Postharvest Losses.



Konstantinos G. Batziakas, Helena Pontes Chiebao, Cary L. Rivard and Eleni D. Pliakoni


Department of Horticulture & Natural Resources, Kansas State University, USA

Materials & Methods Pre-harvest Study

K- State Olathe Horticulture Research and Extension Center, Olathe Kansas




- 2014-2016
- 6 replicates, RCBD
- 6 plants/replicate
 - 'BHN 589'
 - 'Cherokee Purple'




Tomato Yield


2014-2016 Yield per growing system ($p < 0.01$)



PAR (P < 0.0001)



PAR (P < 0.0001)



Tomato – Nutritional Quality

Quality at day of harvest ($p < 0.01$)

BHN589	High Tunnel	Open Field
ORAC	743.5 (64.1)A	723.0 (40.8)A
FRAP	295.9 (21.6)A	315.1 (25.0)A
Total Phenolics	18.6 (2.60)A	21.2 (4.28)A
Sugar/Acid	11.5 (1.27)A	9.9 (0.87)A
Lycopene	4.6 (0.48)A	4.8 (0.66)A
β -Carotene	2.8 (0.17)A	2.6 (0.11)A
Cherokee Purple	High Tunnel	Open Field
ORAC	535.4 (38.6)A	775.2 (103.9)B
FRAP	158.0 (23.6) A	300.5 (50.4) B
Total Phenolics	22.3 (0.51)A	22.1 (2.55)A
Sugar/Acid	13.3 (1.97)A	13.3 (1.18)A
Lycopene	4.1 (0.28)A	5.5 (0.90)A
β -Carotene	1.8 (0.44)A	2.5 (0.31)A

Spinach – Nutritional Quality

Growing Season	Treatment	<u>Total Phenolic content</u> (GA eq/100g FW)	<u>FRAP</u> (μ Mol TE/100g FW)	<u>ORAC</u> (μ Mol TE/100g FW)	<u>Chlorophyll content</u> (mg/100g FW)	<u>Ascorbic Acid content</u> (μ gAA/g FW)
2 (2015/2016)	High Tunnel	65.1 a^z (12.4)	145.3 b (30.3)	3676.9 b (636.6)	100.2 a (26.9)	672.3 b (115.1)
	Open Field	70.5 a (16.6)	225.1 a (52.8)	4888.9 a (597.9)	96.6 a (19.7)	759.7 a (135.9)
	P value	NS	***	***	NS	*
3 (2016/2017)	High Tunnel	49.7 b (9.4)	230.2 b (51.5)	3707.9 b (1166.1)	230.2 b (51.5)	786.7 a (160.3)
	Open Field	70.8 a (14.9)	301.8 a (102.7)	4442.1 a (638.3)	301.8 a (102.7)	838.3 a (205.7)
	P value	***	***	*	***	NS

- Antioxidant capacity decreased in HT
- Vitamin C and Chlorophyll inconsistent
- Impact of environment

Diffuse Poly

Power of Refraction

- Light is refracted as it enters the tunnel
- Trade names
 - Luminance
 - Polar Ice
- ~85% light transmission (reportedly)
- Can reduce temperatures



Diffuse



Standard








PAR (P < 0.0001)



Soil Temperature - Lettuce

Code	Treatment
SP	Std. poly
SP2W	Std. poly, removal 2 weeks prior to harvest
DP	Diffuse poly
CP	Clear poly
UVAUVB	UVA, UVB blocking
SCSP	55% shade cloth, std. poly
OF	Open field




Shade Cloth

So Much to Think About

- New area of research
- Few recommendations
- Timing of application
- White vs. Black vs. Metal
 - Depends on the goal
- Disadvantages:
 - Crop growth
 - Cost



PAR (P < 0.0001)



Diffuse

Clear

UVA/
UVB

55%
Shade

c







Light effects during pre- and postharvest stages to improve nutrition of high tunnel vegetables

Kelly Gude, Ph.D. student, Horticulture and Natural Resources






OBJECTIVE: Evaluate the impact of different poly films on spectral quality and a simulated movable tunnel on light intensity regarding photosynthesis, temperature, crop yield, storage life, and nutritional quality of tomato and lettuce.








Temperature – Tomato

TRT - TOMATO	MAX - Canopy		MIN - Canopy		MAX - Soil		MIN -Soil	
STANDARD	29.8	bcd	19.5	b	25.3	bc	23.8	
REMOVAL	29.2	cd	21.2	a	25.5	b	23.8	
DIFFUSE	28.8	d	19.8	b	24.8	c	23.7	
CLEAR	30.3	abc	19.5	b	26.4	a	24.3	
BLOCK	30.9	a	19.4	b	26.2	a	23.8	
SHADE	30.3	ab	20.3	ab	25.8	ab	23.9	
OPEN	31.5	a	20.0	b	26.4	a	24.1	
	<.0001		<.0001		<.0001		NS	




Photosynthesis and PAR – Tomato



Combined Yield – Tomato

	Total Yield (kg/plant)	Marketable Yield (kg/plant)	Total Fruit Weight (kg/fruit)	Marketable Fruit Weight (kg/fruit)	Marketabili ty (% by weight)	Plant Biomass (g)
Standard	6.53 A ²	5.21 A	0.18 A	0.19 ab	76.5 ab	310
Removal	6.17 A	4.53 AB	0.18 A	0.18 ab	69.5 b	248
Diffuse	7.17 A	5.67 A	0.18 A	0.18 ab	73.3 ab	246
Clear	7.39 A	5.62 A	0.19 A	0.19 a	70.6 ab	278
Block	6.62 A	5.13 A	0.18 A	0.19 a	73.9 ab	311
Shade	4.40 B	3.72 B	0.15 B	0.15 b	77.7 a	209
P-value	<.001	<.001	<.001	< 0.05	< 0.05	NS ³




Combined Yield – Tomato

	Total Yield (kg/plant)	Marketable Yield (kg/plant)	Total Fruit Weight (kg/fruit)	Marketable Fruit Weight (kg/fruit)	Marketabili ty (% by weight)	Plant Biomass (g)
Standard	6.53 A ²	5.21 A	0.18 A	0.19 ab	76.5 ab	310
Removal	6.17 A	4.53 AB	0.18 A	0.18 ab	69.5 b	248
Diffuse	7.17 A	5.67 A	0.18 A	0.18 ab	73.3 ab	246
Clear	7.39 A	5.62 A	0.19 A	0.19 a	70.6 ab	278
Block	6.62 A	5.13 A	0.18 A	0.19 a	73.9 ab	311
Shade	4.40 B	3.72 B	0.15 B	0.15 b	77.7 a	209
P-value	<.001	<.001	<.001	< 0.05	< 0.05	NS ³




Combined Yield – Tomato

	Total Yield (kg/plant)	Marketable Yield (kg/plant)	Total Fruit Weight (kg/fruit)	Marketable Fruit Weight (kg/fruit)	Marketabili ty (% by weight)	Plant Biomass (g)
Standard	6.53 A	5.21 A	0.18 A	0.19 ab	76.5 ab	310
Removal	6.17 A	4.53 AB	0.18 A	0.18 ab	69.5 b	248
Diffuse	7.17 A	5.67 A	0.18 A	0.18 ab	73.3 ab	246
Clear	7.39 A	5.62 A	0.19 A	0.19 a	70.6 ab	278
Block	6.62 A	5.13 A	0.18 A	0.19 a	73.9 ab	311
Shade	4.40 B	3.72 B	0.15 B	0.15 b	77.7 a	209
P-value	<.001	<.001	<.001	< 0.05	< 0.05	NS ³

Combined Yield – Tomato

	Total Yield (kg/plant)	Marketable Yield (kg/plant)	Total Fruit Weight (kg/fruit)	Marketable Fruit Weight (kg/fruit)	Marketabili ty (% by weight)	Plant Biomass (g)
Standard	6.53 A	5.21 A	0.18 A	0.19 ab	76.5 ab	310
Removal	6.17 A	4.53 AB	0.18 A	0.18 ab	69.5 b	248
Diffuse	7.17 A	5.67 A	0.18 A	0.18 ab	73.3 ab	246
Clear	7.39 A	5.62 A	0.19 A	0.19 a	70.6 ab	278
Block	6.62 A	5.13 A	0.18 A	0.19 a	73.9 ab	311
Shade	4.40 B	3.72 B	0.15 B	0.15 b	77.7 a	209
P-value	<.001	<.001	<.001	< 0.05	< 0.05	NS ³



Vitamin C – Tomato





Figure 3. Ascorbic acid content (SE) in tomatoes harvested at 'breaker' and 'light red' maturity once reaching 'mature red' maturity (BR_MR and LR_MR, respectively). Significance: * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$.

Individual Phenolics – Tomato


QUERCETIN

■ BR_MR
▨ LR_MR




Caffeic acid o hexoside

■ BR_MR*
▨ LR_MR




RUTIN

■ BR_MR
▨ LR_MR



U 254

■ BR_MR
▨ LR_MR





PAR – Lettuce

Trts			PAR		
	Fall, Two Star	Spring, Two Star	Fall, Red Fire	Spring, Red Fire	
Standard	347.831	1340.78 a	503.105 a	1465.37	a
Removal	380.15	1498.06 a	614.634 a	1685.3	a
Diffuse	361.391	947.11 ab	373.17 ab	1139.28	ab
Clear	317.033	1278.78 a	496.098 a	1189.26	ab
Block	334.297	1050.73 ab	505.566 a	1416.71	a
Shade	171.547	474.42 b	218.683 b	562.29	b
P-value	ns	0.0076	0.0013	0.017	

Photosynthesis – Lettuce

Trts	Photosynthesis Rate (Pn)						
	Fall, Two Star		Spring, Two Star	Fall, Red Fire		Spring, Red Fire	
Standard	5.20		26.66	4.74		18.28	
Removal	4.08		23.22	5.18		12.11	
Diffuse	4.13		12.41	3.88		13.86	
Clear	3.00		21.17	4.49		18.59	
Block	3.17		21.58	4.94		18.32	
Shade	5.03		14.84	2.81		14.37	
P-value	ns		ns	ns		ns	

Yield – Green Leaf Lettuce

Yield analysis from fall harvests in 2017 (11/3-11/17) & 2018 (12/7-12/13) and spring harvests in 2018 (5/3-5/10).

*OF not statistically analyzed

Code	Treatment	Green Leaf					
		Trt	Fresh Wt. (g/plant)	Core length (cm)	Leaf Area (cm²)	Moisture Loss (%)	DM
SP	Std. poly						
SP2W	Std. poly, removal 2 weeks prior to harvest						
DP	Diffuse poly	SP	68.67 A	18.43	127.22	18.23 AB	8.92
CP	Clear poly						
UVAUVB	UVA, UVB blocking	SP2W	79.67 A	18.13	139.27	15.97 B	8.82
SCSP	55% shade cloth, std. poly	DP	68.69 A	18.31	142.4	19.39 AB	8.60
OF	Open field	CP	62.44 AB	17.48	136.07	18.78 AB	8.64
		UVAUVB	56.65 AB	16.67	124.92	19.49 AB	8.68
		SCSP	40.68 B	17.52	120.06	25.5 A	8.14
		*OF	17.51				
		P	< .01	ns	ns	< .1	ns

Yield – Green Leaf Lettuce

Yield analysis from fall harvests in 2017 (11/3-11/17) & 2018 (12/7-12/13) and spring harvests in 2018 (5/3-5/10).


*OF not statistically analyzed

Code	Treatment	Red Leaf					
		Trt	Fresh Wt. (g/plant)	Core length (cm)	Leaf Area (cm ²)	Moisture Loss (%)	DM
SP	Std. poly	SP	51.93 A	17.26 A	141.5 A	11.48	7.71
SP2W	Std. poly, removal 2 weeks prior to harvest	SP2W	51 A	16.06 ABC	120.28 AB	11.53	7.56
DP	Diffuse poly	DP	52.13 A	16.71 ABC	120.59 AB	11.17	7.27
CP	Clear poly	CP	45.57 AB	15.58 BC	105.32 AB	11.21	7.22
UVAUV	UVA, UVB	UVAUV	39.58 AB	15.16 C	96.04 B	15.04	7.77
B	blocking	B	29.61 B	16.18 ABC	99.1 B	16.56	7.91
SCSP	55% shade cloth, std. poly	SCSP	14.1				
OF	Open field	*OF					
		P	< .01	< .05	ns	ns	




Color - Lettuce

Effects Table	'Two Star'		'Red Fire'	
	Hue	Chroma	Hue	Chroma
SEASON	<.0001	0.0393	<.0001	<.0001
TRT	0.0002	0.0003	<.0001	<.0001
SEASON*TRT	0.0011	0.0003	0.5687	0.5918
DAY	0.1224	0.0088	0.1901	0.0129
SEASON*DAY	0.0035	<.0001	0.7033	0.0758
TRT*DAY	0.0367	0.2126	0.9225	0.9957
SEASON*TRT*DAY	0.0655	0.0299	0.853	0.9032




Phenols – Red Leaf Lettuce


CHICORIC ACID 'Red Fire'




ISOQuercetin 'Red Fire'



CHLOROGENIC ACID 'Red Fire'




caffeoylmalic acid 'Red Fire'




Phenols – Green Leaf Lettuce


CHICORIC ACID 'Two Star'




ISOQUERCETIN 'Two Star'




CHLOROGENIC ACID 'Two Star'




caffeoylmalic acid 'Two Star'




Consumer Sensory Analysis - Lettuce



Consumer Sensory Analysis - Lettuce



Movable Tunnel Project











Movable Tunnel – Lettuce Yield

Red Leaf					
TRTs	Fresh Wt. (g/head)	Core length (cm)	Leaf Area (cm ²)	Moisture Loss (%)	Dry Matter (%)
SP	76.69	15.90a	140.20A	14.65	7.34A
SP48h	88.11	14.98ab	107.14C	25.90	5.82 B
SP1W	97.36	15.19ab	127.63AB	29.56	5.72 B
SP2W	67.71	14.41b	111.75BC	29.39	6.27 AB

P

<.1

< .01

< .01

Movable Tunnel – Lettuce Yield

Green Leaf					
TRTs	Fresh Wt. (g/head)	Core Length (cm)	Leaf Area (cm ²)	Moisture Loss (%)	Dry Matter (%)
SP	105.97A	17.31A	120.30	18.23	7.60AB
SP48h	122.96A	17.98A	114.88	19.52	6.26B
SP1W	103.05AB	16.83A	103.55	19.31	7.31AB
SP2W	69.60B	13.96B	82.87	17.16	8.47A

P

$< .05$


< .05

Summary

- Manipulating light and microclimate is the next frontier in food production technology
 - Role of HTs
- Research is thin right now, but a few findings from our work:
 - UV light probably important, but not for everything
 - Phytochemicals can behave differently
 - ENVIRONMENT trumps HT covering
 - Light is likely impacting: microclimate, Pn, quality, and yield

Summary

- Grower Recommendations:
 - Standard poly performing well compared to other plastics (including clear)
 - Antioxidants
 - Diffuse plastics comparable with standard
 - Be careful with shade cloth
 - Yield, color (lettuce), phytochemicals
 - Industry recommendations may be suspect (transmission)
 - Movable tunnels provide a NEW option
 - More research is needed (and otw)



Questions?