



TABLE 1: REQUIRED IMMUNIZATIONS – CHILD CARE CENTER/PRESCHOOL*

BY THIS AGE	STUDENTS ARE REQUIRED TO HAVE							
	DTAP (Diphtheria, Tetanus, Pertussis)	POLIO	HIB** (<i>Haemophilus influenzae</i> type B)	PCV** (Pneumococcal Conjugate Vaccine)	HEPATITIS B	HEPATITIS A	MMR (Measles, Mumps, Rubella)	VARICELLA (Chickenpox)
3 MONTHS	1 dose	1 dose	1 dose	1 dose	2 doses	Not required at this age	Not required at this age	Not required at this age
5 MONTHS	2 doses	2 doses	2 doses	2 doses	2 doses	Not required at this age	Not required at this age	Not required at this age
7 MONTHS	3 doses	2 doses	2 - 3 doses [†]	3 doses	2 doses	Not required at this age	Not required at this age	Not required at this age
16 MONTHS	3 doses	2 doses	3 - 4 doses [¶]	4 doses [§]	2 doses	Not required at this age	1 dose	1 dose
19 MONTHS	4 doses	3 doses	3 - 4 doses [¶]	4 doses [§]	3 doses	1 dose	1 dose	1 dose
24 MONTHS	4 doses	3 doses	3 - 4 doses [¶]	4 doses [§]	3 doses	2 doses	1 dose	1 dose

*SEE TABLE 6: EXCEPTIONS TO IMMUNIZATION REQUIREMENTS

**FOR INFORMATION ON CHILDREN WHO START THE HIB OR PCV SERIES AT AGE 7 MONTHS OR OLDER, SEE TABLE 5A AND 5B

[†]If all PedvaxHIB® (Merck) administered, 2 doses are required

If ActHIB® (sanofi), Pentacel® (sanofi), or Hiberix® (GSK) administered, 3 doses are required

If both PedvaxHIB® and either ActHIB®, Pentacel®, or Hiberix® administered, or if product type is UNKNOWN or not listed here, 3 doses are required

[¶]If all PedvaxHIB® (Merck) administered at the routinely recommended ages (2, 4, 12-15 months), 3 doses are required.

If ActHIB® (sanofi), Pentacel® (sanofi), or Hiberix® (GSK) administered at the routinely recommended ages (2, 4, 6, 12-15 months), 4 doses are required.

If both PedvaxHIB® and either ActHIB®, Pentacel®, or Hiberix® administered at the routinely recommended ages (2, 4, 6, 12-15 months), or if product type is UNKNOWN or not listed here, 4 doses are required.

The number of doses required for children who did not receive Hib at the routinely recommended ages depends on the child’s age at the time of administration of each dose and the vaccine product. See Table 5A.

[§]If administered at the routinely recommended ages (2, 4, 6, 12-15 months), 4 doses of PCV are required.

The number of doses required for children who did not receive PCV at the routinely recommended ages depends on the child’s age at the time of administration of each dose. See Table 5B.